

QUARK MASS HIERARCHY AND FLAVOR MIXING

Harald Fritzsch

Sektion Physik, Universität München,
D-80333 München, Germany

ABSTRACT

In view of the observed strong hierarchy of the quark and lepton masses and of the flavor mixing angles it is argued that the description of flavor mixing must take this into account. One particular interesting way to describe the flavor mixing, which, however, is not the one used today, emerges, which is best suited for models of quark mass matrices based on flavor symmetries. We conclude that the unitarity triangle important for B physics should be close to or identical to a rectangular triangle. CP violation is maximal in this sense.

*Invited talk given at the
International Conference on Orbis Scientiae 1997:
Physics of Mass
Miami Beach, Florida (December 12–15, 1997)*

The phenomenon of flavor mixing, which is intrinsically linked to CP -violation, is an important ingredient of the Standard Model of Basic Interactions. Yet unlike other features of the Standard Model, e. g. the mixing of the neutral electroweak gauge bosons, it is a phenomenon which can merely be described. A deeper understanding is still lacking, but most theoreticians would agree that it is directly linked to the mass spectrum of the quarks – the possible mixing of lepton flavors will not be discussed here. Furthermore there is a general consensus that a deeper dynamical understanding would require to go beyond the physics of the Standard Model. In this talk I shall not go thus far. Instead I shall demonstrate that the observed properties of the flavor mixing, combined with our knowledge about the quark mass spectrum, suggest specific symmetry properties which allow to fix the flavor mixing parameters with high precision, thus predicting the outcome of the experiments which will soon be performed at the B -meson factories.

In the standard electroweak theory, the phenomenon of flavor mixing of the quarks is described by a 3×3 unitary matrix, the Cabibbo-Kobayashi-Maskawa (CKM) matrix^{1,2}. This matrix can be expressed in terms of four parameters, which are usually taken as three rotation angles and one phase. A number of different parametrizations have been proposed in the literature^{2,3,4,5}. Of course, adopting a particular parametrization of flavor mixing is arbitrary and not directly a physics issue. Nevertheless it is quite likely that the actual values of flavor mixing parameters (including the strength of CP violation), once they are known with high precision, will give interesting information about the physics beyond the standard model. Probably at this point it will turn out that a particular description of the CKM matrix is more useful and transparent than the others. For this reason, let me first analyze all possible parametrizations and point out their respective advantages and disadvantages.

In the standard model the quark flavor mixing arises once the up- and down-type mass matrices are diagonalized. The generation of quark masses is intimately related to the phenomenon of flavor mixing. In particular, the flavor mixing parameters do depend on the elements of quark mass matrices. A particular structure of the underlying mass matrices calls for a particular choice of the parametrization of the flavor mixing matrix. For example, in it was noticed⁶ that a rather special form of the flavor mixing matrix results, if one starts from Hermitian mass matrices in which the (1,3) and (3,1) elements vanish. This has been subsequently observed again in a number of papers⁷. Recently we have studied the exact form of such a description from a general point of view and pointed out many advantages of this type of representation in the discussion of flavor mixing and CP -violating phenomena⁵, which will be discussed later.

In the standard model the weak charged currents are given by

$$\overline{(u, c, t)}_L \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}_L, \quad (1)$$

where $u, c, \dots, b$ are the quark mass eigenstates, L denotes the left-handed fields, and V_{ij} are elements of the CKM matrix V . In general V_{ij} are complex numbers, but their absolute values are measurable quantities. For example, $|V_{cb}|$ primarily determines the lifetime of B mesons. The phases of V_{ij} , however, are not physical, like the phases of quark fields. A phase transformation of the u quark ($u \rightarrow u e^{i\alpha}$), for example, leaves the quark mass term invariant but changes the elements in the first row of V (i.e., $V_{uj} \rightarrow V_{uj} e^{-i\alpha}$). Only a common phase transformation of all quark fields leaves all elements of V invariant, thus there is a five-fold freedom to adjust the phases of V_{ij} .

In general the unitary matrix V depends on nine parameters. Note that in the absence of complex phases V would consist of only three independent parameters, corresponding to three (Euler) rotation angles. Hence one can describe the complex matrix V by three angles and six phases. Due to the freedom in redefining the quark field phases, five of the six phases in V can be absorbed, and we arrive at the well-known result that the CKM matrix V can be parametrized in terms of three rotation angles and one CP -violating phase. The question about how many different ways to describe V may exist was raised some time ago⁸. Recently the problem was reconsidered and brought in connection with the mass hierarchy⁵.

In our view the best possibility to describe the flavor mixing in the standard model is to adopt the parametrization discussed in ref. (5). This parametrization has a number of significant advantages. In the following part I shall show that this parametrization follows automatically if we impose the constraints from the chiral symmetries and the hierarchical structure of the mass eigenvalues^{9,10,11}. We take the point of view that the quark mass eigenvalues are dynamical entities, and one could change their values in order to study certain symmetry limits, as it is done in QCD. In the standard electroweak model, in which the quark mass matrices are given by the coupling of a scalar field to various quark fields, this can certainly be done by adjusting the related coupling constants. Whether it is possible in reality is an open question. It is well-known that the quark mass matrices can always be made hermitian by a suitable transformation of the right-handed fields. Without loss of generality, we shall suppose in this paper that the quark mass matrices are hermitian. In the limit where the masses of the u and d quarks are set to zero, the quark mass matrix $\tilde{M}$ (for both charge $+2/3$ and charge $-1/3$ sectors) can be arranged such that its elements $\tilde{M}_{i1}$ and $\tilde{M}_{1i}$ ($i = 1, 2, 3$) are all zero^{9,10}. Thus the

quark mass matrices have the form

$$\tilde{M} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \tilde{C} & \tilde{B} \\ 0 & \tilde{B}^* & \tilde{A} \end{pmatrix}. \quad (2)$$

The observed mass hierarchy is incorporated into this structure by denoting the entry which is of the order of the t -quark or b -quark mass by $\tilde{A}$, with $\tilde{A} \gg \tilde{C}, |\tilde{B}|$. It can easily be seen (see, e.g., ref. (12)) that the complex phases in the mass matrices (1) can be rotated away by subjecting both $\tilde{M}_u$ and $\tilde{M}_d$ to the same unitary transformation. Thus we shall take $\tilde{B}$ to be real for both up- and down-quark sectors. As expected, CP violation cannot arise at this stage. The diagonalization of the mass matrices leads to a mixing between the second and third families, described by an angle $\tilde{\theta}$. The flavor mixing matrix is then given by

$$\tilde{V} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \tilde{c} & \tilde{s} \\ 0 & -\tilde{s} & \tilde{c} \end{pmatrix}, \quad (3)$$

where $\tilde{s} \equiv \sin \tilde{\theta}$ and $\tilde{c} \equiv \cos \tilde{\theta}$. In view of the fact that the limit $m_u = m_d = 0$ is not far from reality, the angle $\tilde{\theta}$ is essentially given by the observed value of $|V_{cb}|$ ($= 0.039 \pm 0.002$)^{13,14}; i.e., $\tilde{\theta} = 2.24^\circ \pm 0.12^\circ$.

At the next and final stage of the chiral evolution of the mass matrices, the masses of the u and d quarks are introduced. The Hermitian mass matrices have in general the form:

$$M = \begin{pmatrix} E & D & F \\ D^* & C & B \\ F^* & B^* & A \end{pmatrix} \quad (4)$$

with $A \gg C, |B| \gg E, |D|, |F|$. By a common unitary transformation of the up- and down-type quark fields, one can always arrange the mass matrices M_u and M_d in such a way that $F_u = F_d = 0$; i.e.,

$$M = \begin{pmatrix} E & D & 0 \\ D^* & C & B \\ 0 & B^* & A \end{pmatrix}. \quad (5)$$

This can easily be seen as follows. If phases are neglected, the two symmetric mass matrices M_u and M_d can be transformed by an orthogonal transformation matrix O , which can be described by three angles such that they assume the form (5). The condition $F_u = F_d = 0$ gives two constraints for the three angles of O . If complex phases are allowed in M_u and M_d , the condition $F_u = F_u^* = F_d = F_d^* = 0$ imposes four constraints, which can also be fulfilled, if M_u and M_d are subjected to a common unitary transformation matrix U . The latter depends on nine parameters. Three of them are not suitable for our purpose, since they are just diagonal phases; but the

remaining six can be chosen such that the vanishing of F_u and F_d results.

The basis in which the mass matrices take the form (5) is a basis in the space of quark flavors, which in our view is of special interest. It is a basis in which the mass matrices exhibit two texture zeros, for both up- and down-type quark sectors. These, however, do not imply special relations among mass eigenvalues and flavor mixing parameters (as pointed out above). In this basis the mixing is of the “nearest neighbour” form, since the (1,3) and (3,1) elements of M_u and M_d vanish; no direct mixing between the heavy t (or b) quark and the light u (or d) quark is present (see also ref. (15)). In certain models (see, e.g., ref. (16)), this basis is indeed of particular interest, but we shall proceed without relying on a special texture models for the mass matrices.

A mass matrix of the type (5) can in the absence of complex phases be diagonalized by a rotation matrix, described by two angles only. At first the off-diagonal element B is rotated away by a rotation between the second and third families (angle θ_{23}); at the second step the element D is rotated away by a transformation of the first and second families (angle θ_{12}). No rotation between the first and third families is required. The rotation matrix for this sequence takes the form

$$\begin{aligned} R &= R_{12}R_{23} = \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \\ &= \begin{pmatrix} c_{12} & s_{12}c_{23} & s_{12}s_{23} \\ -s_{12} & c_{12}c_{23} & c_{12}s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix}, \end{aligned} \quad (6)$$

where $c_{12} \equiv \cos \theta_{12}$, $s_{12} \equiv \sin \theta_{12}$, etc. The flavor mixing matrix V is the product of two such matrices, one describing the rotation among the up-type quarks, and the other describing the rotation among the down-type quarks:

$$V = R_{12}^u R_{23}^u (R_{23}^d)^{-1} (R_{12}^d)^{-1}. \quad (7)$$

The product $R_{23}^u (R_{23}^d)^{-1}$ can be written as a rotation matrix described by a single angle θ . In the limit $m_u = m_d = 0$, this is just the angle $\tilde{\theta}$ encountered in eq. (6). The angle which describes the R_{12}^u rotation shall be denoted by θ_u ; the corresponding angle for the R_{12}^d rotation by θ_d . Thus in the absence of CP -violating phases the flavor mixing matrix takes the following specific form:

$$\begin{aligned} V &= \begin{pmatrix} c_u & s_u & 0 \\ -s_u & c_u & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & c & s \\ 0 & -s & c \end{pmatrix} \begin{pmatrix} c_d & -s_d & 0 \\ s_d & c_d & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} s_u s_d c + c_u c_d & s_u c_d c - c_u s_d & s_u s \\ c_u s_d c - s_u c_d & c_u c_d c + s_u s_d & c_u s \\ -s_d s & -c_d s & c \end{pmatrix}, \end{aligned} \quad (8)$$

where $c_u \equiv \cos \theta_u$, $s_u \equiv \sin \theta_u$, etc.

We proceed by including the phase parameters of the quark mass matrices in eq. (5). It can easily be seen that, by suitable rephasing of the quark fields, the flavor mixing matrix can finally be written in terms of only a single phase φ as follows:

$$\begin{aligned} V &= \begin{pmatrix} c_u & s_u & 0 \\ -s_u & c_u & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} e^{-i\varphi} & 0 & 0 \\ 0 & c & s \\ 0 & -s & c \end{pmatrix} \begin{pmatrix} c_d & -s_d & 0 \\ s_d & c_d & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} s_u s_d c + c_u c_d e^{-i\varphi} & s_u c_d c - c_u s_d e^{-i\varphi} & s_u s \\ c_u s_d c - s_u c_d e^{-i\varphi} & c_u c_d c + s_u s_d e^{-i\varphi} & c_u s \\ -s_d s & -c_d s & c \end{pmatrix}. \end{aligned} \quad (9)$$

Note that the three angles θ_u , θ_d and θ in eq. (12) can all be arranged to lie in the first quadrant through a suitable redefinition of quark field phases. Consequently all s_u , s_d , s and c_u , c_d , c are positive. The phase φ can in general take values from 0 to 2π ; and CP violation is present in the weak interactions if $\varphi \neq 0, \pi$ and 2π .

This particular representation of the flavor mixing matrix is the main result of this paper. In comparison with all other parametrizations discussed previously^{2,3}, it has a number of interesting features which in our view make it very attractive and provide strong arguments for its use in future discussions of flavor mixing phenomena, in particular, those in B -meson physics (see also refs. (17, 18)). We shall discuss them below.

a) The flavor mixing matrix V in eq. (12) follows directly from the chiral expansion of the mass matrices. Thus it naturally takes into account the hierarchical structure of the quark mass spectrum.

b) The complex phase φ describing CP violation appears only in the (1,1), (1,2), (2,1) and (2,2) elements of V , i.e., in the elements involving only the quarks of the first and second families. This is a natural description of CP violation since in our hierarchical approach CP violation is not directly linked to the third family, but rather to the first and second ones, and in particular to the mass terms of the u and d quarks.

It is instructive to consider the special case $s_u = s_d = s = 0$. Then the flavor mixing matrix V takes the form

$$V = \begin{pmatrix} e^{-i\varphi} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (10)$$

This matrix describes a phase change in the weak transition between u and d , while no phase change is present in the transitions between c and s as well as t and

b. Of course, this effect can be absorbed in a phase change of the u - and d -quark fields, and no CP violation is present. Once the angles θ_u , θ_d and θ are introduced, however, CP violation arises. It is due to a phase change in the weak transition between u' and d' , where u' and d' are the rotated quark fields, obtained by applying the corresponding rotation matrices given in eq. (9) to the quark mass eigenstates (u' : mainly u , small admixture of c ; d' : mainly d , small admixture of s).

Since the mixing matrix elements involving the t or b quarks are real in the representation (9), one can find that the phase parameter of B_q^0 - $\bar{B}_q^0$ mixing ($q = d$ or s), dominated by the box-diagram contributions in the standard model¹⁹, is essentially unity:

$$\left(\frac{q}{p}\right)_{B_q} = \frac{V_{tb}^* V_{tq}}{V_{tb} V_{tq}^*} = 1. \quad (11)$$

In most other parametrizations of the flavor mixing matrix, however, the two rephasing-variant quantities $(q/p)_{B_d}$ and $(q/p)_{B_s}$ take different (maybe complex) values.

c) The dynamics of flavor mixing can easily be interpreted by considering certain limiting cases in eq. (9). In the limit $\theta \rightarrow 0$ (i.e., $s \rightarrow 0$ and $c \rightarrow 1$), the flavor mixing is, of course, just a mixing between the first and second families, described by only one mixing angle (the Cabibbo angle θ_C ¹).

It is a special and essential feature of the representation (9) that the Cabibbo angle is *not* a basic angle, used in the parametrization. The matrix element V_{us} (or V_{cd}) is indeed a superposition of two terms including a phase. This feature arises naturally in our hierarchical approach, but it is not new. In many models of specific textures of mass matrices, it is indeed the case that the Cabibbo-type transition V_{us} (or V_{cd}) is a superposition of several terms. At first, it was obtained by one of the authors in the discussion of the two-family mixing²⁰.

In the limit $\theta = 0$ considered here, one has $|V_{us}| = |V_{cd}| = \sin \theta_C \equiv s_C$ and

$$s_C = \left| s_u c_d - c_u s_d e^{-i\varphi} \right|. \quad (12)$$

This relation describes a triangle in the complex plane, as illustrated in Fig. 1, which we shall denote as the “LQ- triangle” (“light quark triangle”). This triangle is a feature of the mixing of the first two families (see also ref. (20)). Explicitly one has (for $s = 0$):

$$\tan \theta_C = \sqrt{\frac{\tan^2 \theta_u + \tan^2 \theta_d - 2 \tan \theta_u \tan \theta_d \cos \varphi}{1 + \tan^2 \theta_u \tan^2 \theta_d + 2 \tan \theta_u \tan \theta_d \cos \varphi}}. \quad (13)$$

Certainly the flavor mixing matrix V cannot accommodate CP violation in this limit. However, the existence of φ seems necessary in order to make eq. (16)




Abbildung 1: The LQ-triangle in the complex plane.

compatible with current data, as one can see below.

d) The three mixing angles θ , θ_u and θ_d have a precise physical meaning. The angle θ describes the mixing between the second and third families, which is generated by the off-diagonal terms B_u and B_d in the up and down mass matrices of eq. (9). We shall refer to this mixing involving t and b as the “heavy quark mixing”. The angle θ_u , however, primarily describes the u - c mixing, corresponding to the D_u term in M_u . We shall denote this as the “u-channel mixing”. The angle θ_d primarily describes the d - s mixing, corresponding to the D_d term in M_d ; this will be denoted as the “d-channel mixing”. Thus there exists an asymmetry between the mixing of the first and second families and that of the second and third families, which in our view reflects interesting details of the underlying dynamics of flavor mixing. The heavy quark mixing is a combined effect, involving both charge $+2/3$ and charge $-1/3$ quarks, while the u- or d-channel mixing (described by the angle θ_u or θ_d) proceeds solely in the charge $+2/3$ or charge $-1/3$ sector. Therefore an experimental determination of these two angles would allow to draw interesting conclusions about the amount and perhaps the underlying pattern of the u- or d-channel mixing.

e) The three angles θ , θ_u and θ_d are related in a very simple way to observable quantities of B -meson physics.

For example, θ is related to the rate of the semileptonic decay $B \rightarrow D^* l \nu_l$; θ_u is associated with the ratio of the decay rate of $B \rightarrow (\pi, \rho) l \nu_l$ to that of $B \rightarrow D^* l \nu_l$; and θ_d can be determined from the ratio of the mass difference between two B_d mass eigenstates to that between two B_s mass eigenstates. From eq. (9) we find the following exact relations:

$$\sin \theta = |V_{cb}| \sqrt{1 + \left| \frac{V_{ub}}{V_{cb}} \right|^2}, \quad (14)$$

and

$$\begin{aligned} \tan \theta_u &= \left| \frac{V_{ub}}{V_{cb}} \right|, \\ \tan \theta_d &= \left| \frac{V_{td}}{V_{ts}} \right|. \end{aligned} \quad (15)$$

These simple results make the parametrization (9) uniquely favorable for the study of B -meson physics.

By use of current data on $|V_{ub}|$ and $|V_{cb}|$, i.e., $|V_{cb}| = 0.039 \pm 0.002^{13,14}$ and $|V_{ub}/V_{cb}| = 0.08 \pm 0.02^{14}$, we obtain $\theta_u = 4.57^\circ \pm 1.14^\circ$ and $\theta = 2.25^\circ \pm 0.12^\circ$. Taking $|V_{td}| = (8.6 \pm 2.1) \times 10^{-314}$, which was obtained from the analysis of current data on B_d^0 - $\bar{B}_d^0$ mixing, we get $|V_{td}/V_{ts}| = 0.22 \pm 0.07$, i.e., $\theta_d = 12.7^\circ \pm 3.8^\circ$. Both the heavy quark mixing angle θ and the u-channel mixing angle θ_u are relatively small. The smallness of θ implies that Eqs. (11) and (12) are valid to a high degree of precision (of order $1 - c \approx 0.001$).

f) According to eq. (12), as well as eq. (11), the phase φ is a phase difference between the contributions to V_{us} (or V_{cd}) from the u-channel mixing and the d-channel mixing. Therefore φ is given by the relative phase of D_d and D_u in the quark mass matrices (4), if the phases of B_u and B_d are absent or negligible.

The phase φ is not likely to be 0° or 180° , according to the experimental values given above, even though the measurement of CP violation in K^0 - $\bar{K}^0$ mixing¹⁹ is not taken into account. For $\varphi = 0^\circ$, one finds $\tan \theta_C = 0.14 \pm 0.08$; and for $\varphi = 180^\circ$, one gets $\tan \theta_C = 0.30 \pm 0.08$. Both cases are barely consistent with the value of $\tan \theta_C$ obtained from experiments ($\tan \theta_C \approx |V_{us}/V_{ud}| \approx 0.226$).

g) The CP -violating phase φ in the flavor mixing matrix V can be determined from $|V_{us}|$ ($= 0.2205 \pm 0.0018$)¹⁹ through the following formula, obtained easily from eq. (8):

$$\varphi = \arccos \left(\frac{s_u^2 c_d^2 c^2 + c_u^2 s_d^2 - |V_{us}|^2}{2s_u c_u s_d c_d c} \right). \quad (16)$$

The two-fold ambiguity associated with the value of φ , coming from $\cos \varphi = \cos(2\pi - \varphi)$, is removed if one takes $\sin \varphi > 0$ into account (this is required by current data on CP violation in K^0 - $\bar{K}^0$ mixing (i.e., ϵ_K). More precise measurements of the angles θ_u and θ_d in the forthcoming experiments of B physics will reduce the uncertainty of φ to be determined from eq. (19). This approach is of course complementary to the direct determination of φ from CP asymmetries in some weak B -meson decays into hadronic CP eigenstates²¹.

Considering the presently known phenomenological constraints (see e.g. Ref.²²) the value of φ is most likely in the range 40° to 120° ; the central value is $\varphi \approx 81^\circ$. Note that φ is essentially independent of the angle θ , due to the tiny observed value of the latter. Once $\tan \theta_d$ is precisely measured, one shall be able to fix the magnitude of φ to a satisfactory degree of accuracy.

h) It is well-known that CP violation in the flavor mixing matrix V can be rephasing-invariantly described by a universal quantity $\mathcal{J}^{23}$:

$$\text{Im} \left(V_{il} V_{jm} V_{im}^* V_{jl}^* \right) = \mathcal{J} \sum_{k,n=1}^3 (\epsilon_{ijk} \epsilon_{lmn}) . \quad (17)$$

In the parametrisation (9), $\mathcal{J}$ reads

$$\mathcal{J} = s_u c_u s_d c_d s^2 c \sin \varphi \quad (18)$$

Obviously $\varphi = 90^\circ$ leads to the maximal value of $\mathcal{J}$. Indeed $\varphi = 90^\circ$, a particularly interesting case for CP violation, is quite consistent with current data. In this case the mixing term D_d in eq. (5) can be taken to be real, and the term D_u to be imaginary, if $\text{Im}(B_u) = \text{Im}(B_d) = 0$ is assumed. Since in our description of the flavor mixing the complex phase φ is related in a simple way to the phases of the quark mass terms, the case $\varphi = 90^\circ$ is especially interesting. It can hardly be an accident, and this case should be studied further. The possibility that the phase φ describing CP violation in the standard model is given by the algebraic number $\pi/2$ should be taken seriously. It may provide a useful clue towards a deeper understanding of the origin of CP violation and of the dynamical origin of the fermion masses.

In ref. (5) the case $\varphi = 90^\circ$ has been denoted as “maximal” CP violation. It implies in our framework that in the complex plane the u-channel and d-channel mixings are perpendicular to each other. In this special case (as well as $\theta \rightarrow 0$), we have

$$\tan^2 \theta_C = \frac{\tan^2 \theta_u + \tan^2 \theta_d}{1 + \tan^2 \theta_u \tan^2 \theta_d} . \quad (19)$$

To a good approximation (with the relative error $\sim 2\%$), one finds $s_C^2 \approx s_u^2 + s_d^2$.

i) At future B -meson factories, the study of CP violation will concentrate on measurements of the unitarity triangle.

The unitarity triangle (a) and its rescaled counterpart (b) in the complex plane.

$$S_u + S_c + S_t = 0 , \quad (20)$$

where $S_i \equiv V_{id} V_{ib}^*$ in the complex plane (see Fig. 2(a)). The inner angles of this triangle. are denoted as¹⁹.

$$\begin{aligned} \alpha &\equiv \arg(-S_t S_u^*) , \\ \beta &\equiv \arg(-S_c S_t^*) , \\ \gamma &\equiv \arg(-S_u S_c^*) . \end{aligned} \quad (21)$$

In terms of the parameters θ , θ_u , θ_d and φ , we obtain

$$\begin{aligned}\sin(2\alpha) &= \frac{2c_u c_d \sin \varphi (s_u s_d c + c_u c_d \cos \varphi)}{s_u^2 s_d^2 c^2 + c_u^2 c_d^2 + 2s_u c_u s_d c_d c \cos \varphi}, \\ \sin(2\beta) &= \frac{2s_u c_d \sin \varphi (c_u s_d c - s_u c_d \cos \varphi)}{c_u^2 s_d^2 c^2 + s_u^2 c_d^2 - 2s_u c_u s_d c_d c \cos \varphi}.\end{aligned}\quad (22)$$

To an excellent degree of accuracy, one finds $\alpha \approx \varphi$. In order to illustrate how accurate this relation is, let us input the central values of θ , θ_u and θ_d (i.e., $\theta = 2.25^\circ$, $\theta_u = 4.57^\circ$ and $\theta_d = 12.7^\circ$) to eq. (22). Then one arrives at $\varphi - \alpha \approx 1^\circ$ as well as $\sin(2\alpha) \approx 0.34$ and $\sin(2\beta) \approx 0.65$. It is expected that $\sin(2\alpha)$ and $\sin(2\beta)$ will be directly measured from the CP asymmetries in $B_d \rightarrow \pi^+ \pi^-$ and $B_d \rightarrow J/\psi K_S$ modes at a B -meson factory.

Note that the three sides of the unitarity triangle (21) can be rescaled by $|V_{cb}|$. In a very good approximation (with the relative error $\sim 2\%$), one arrives at

$$|S_u| : |S_c| : |S_t| \approx s_u c_d : s_C : s_d. \quad (23)$$

Equivalently, one can obtain

$$s_\alpha : s_\beta : s_\gamma \approx s_C : s_u c_d : s_d, \quad (24)$$

where $s_\alpha \equiv \sin \alpha$, etc. Comparing the unitarity triangle with the LQ-triangle in Fig. 1, we find that they are indeed congruent with each other to a high degree of accuracy. The congruent relation between these two triangles is particularly interesting, since the LQ-triangle is essentially a feature of the physics of the first two quark families, while the unitarity triangle is linked to all three families. In this connection it is of special interest to note that in models which specify the textures of the mass matrices the Cabibbo triangle and hence three inner angles of the unitarity triangle can be fixed by the spectrum of the light quark masses and the CP -violating phase φ .

j) It is worth pointing out that the u-channel and d-channel mixing angles are related to the so-called Wolfenstein parameters²³ in a simple way:

$$\begin{aligned}\tan \theta_u &= \left| \frac{V_{ub}}{V_{cb}} \right| \approx \lambda \sqrt{\rho^2 + \eta^2}, \\ \tan \theta_d &= \left| \frac{V_{td}}{V_{ts}} \right| \approx \lambda \sqrt{(1 - \rho)^2 + \eta^2},\end{aligned}\quad (25)$$

where $\lambda \approx s_C$ measures the magnitude of V_{us} . Note that the CP -violating parameter η is linked to φ through

$$\sin \varphi \approx \frac{\eta}{\sqrt{\rho^2 + \eta^2} \sqrt{(1 - \rho)^2 + \eta^2}} \quad (26)$$

in the lowest-order approximation. Then $\varphi = 90^\circ$ implies $\eta^2 \approx \rho(1 - \rho)$, on the condition $0 < \rho < 1$. In this interesting case, of course, the flavor mixing matrix can fully be described in terms of only three independent parameters.

k) Compared with the standard parametrization of the flavor mixing matrix V the parametrization (9) has an additional advantage: the renormalization-group evolution of V , from the weak scale to an arbitrary high energy scale, is to a very good approximation associated only with the angle θ . This can easily be seen if one keeps the t and b Yukawa couplings only and neglects possible threshold effect in the one-loop renormalization-group equations of the Yukawa matrices²⁴. Thus the parameters θ_u , θ_d and φ are essentially independent of the energy scale, while θ does depend on it and will change if the underlying scale is shifted, say from the weak scale ($\sim 10^2$ GeV) to the grand unified theory scale (of order 10^{16} GeV). In short, the heavy quark mixing is subject to renormalization-group effects; but the u- and d-channel mixings are not, likewise the phase φ describing CP violation and the LQ-triangle as a whole.

We have presented a new description of the flavor mixing phenomenon, which is based on the phenomenological fact that the quark mass spectrum exhibits a clear hierarchy pattern. This leads uniquely to the interpretation of the flavor mixing in terms of a heavy quark mixing, followed by the u-channel and d-channel mixings. The complex phase φ , describing the relative orientation of the u-channel mixing and the d-channel mixing in the complex plane, signifies CP violation, which is a phenomenon primarily linked to the physics of the first two families. The Cabibbo angle is not a basic mixing parameter, but given by a superposition of two terms involving the complex phase φ . The experimental data suggest that the phase φ , which is directly linked to the phases of the quark mass terms, is close to 90° . This opens the possibility to interpret CP violation as a maximal effect, in a similar way as parity violation.

Our description of flavor mixing has many clear advantages compared with other descriptions. We propose that it should be used in the future description of flavor mixing and CP violation, in particular, for the studies of quark mass matrices and B -meson physics.

The description of the flavor mixing phenomenon given above is of special interest if for the U and D channel mixing the quark mass textures discussed first in²⁰ are applied (see also⁵). In that case one finds²⁵ (apart from small corrections)

$$\tan\Theta_d = \sqrt{\frac{m_d}{m_s}} \quad (27)$$

$$\tan\Theta_u = \sqrt{\frac{m_u}{m_c}}.$$

The experimental value for $\tan \Theta_u$ given by the ratio V_{ub}/V_{cb} is in agreement with the observed value for $(m_u/m_c)^{1/2} \approx 0.07$, but the errors for both $(m_u/m_c)^{1/2}$ and V_{ub}/V_{cb} are the same (about 25%). Thus from the underlying texture no new information is obtained.

This is not true for the angle Θ_d , whose experimental value is due to a large uncertainty: $\Theta_d = 12.7^\circ \pm 3.8^\circ$. If Θ_d is given indeed by the square root of the quark mass ratio m_d/m_s , which is known to a high accuracy, we would know Θ_d and therefore all four parameters of the CKM matrix with high precision.

As emphasized in ref. (5), the phase angle φ is very close to 90° , implying that the LQ-triangle and the unitarity triangle are essentially rectangular triangles. In particular the angle β which is likely to be measured soon in the study of the reaction $B^\circ \rightarrow J/\psi K_s^\circ$ is expected to be close to 20° .

It will be very interesting to see whether the angles Θ_d and Θ_u are indeed given by the square roots of the light quark mass ration m_d/m_s and m_u/m_c , which imply that the phase φ is close to or exactly 90° . This would mean that the light quarks play the most important rôle in the dynamics of flavor mixing and CP violation and that a small window has been opened allowing the first view accross the physics landscape beyond the mountain chain of the Standard Model.

REFERENCES

1. N. Cabibbo, *Phys. Rev. Lett.* 10 (1963) 531.
2. M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* 49 (1973) 652.
3. See, e.g., L. Maiani, *in: Proc. 1977 Int. Symp. on Lepton and Photon Interactions at High Energies (DESY, Hamburg, (1977), 867;*
L.L. Chau and W.Y. Keung, *Phys. Rev. Lett.* 53 (1984) 1802;
H. Fritzsch, *Phys. Rev.* D32 (1985) 3058;
H. Harari and M. Leurer, *Phys. Lett.* B181 (1986) 123;
H. Fritzsch and J. Plankl, *Phys. Rev.* D35 (1987) 1732.
4. F.J. Gilman, Kleinknecht, and B. Renk, *Phys. Rev.* D 54 (1996) 94.
5. H. Fritzsch and Z.Z. Xing, *Phys. Lett.* B353 (1995) 114.
6. H. Fritzsch, *Nucl. Phys.* B155 (1979) 189.
7. S. Dimopoulos, L.J. Hall, and S. Raby, *Phys. Rev.* D45 (1992) 4192;
L.J. Hall and A. Rasin, *Phys. Lett.* B315 (1993) 164;
R. Barbieri, L.J. Hall, and A. Romanino, *Phys. Lett.* B401 (1997) 47.

8. C. Jarlskog, in *CP Violation*, edited by C. Jarlskog (World Scientific, 1989), p. 3.
9. H. Fritzsch, *Phys. Lett.* B184 (1987) 391.
10. H. Fritzsch, *Phys. Lett.* B189 (1987) 191.
11. L.J. Hall and S. Weinberg, *Phys. Rev.* D48 (1993) 979.
12. H. Lehmann, C. Newton, and T.T. Wu, *Phys. Lett.* B384 (1996) 249.
13. M. Neubert, *Int. J. Mod. Phys.* A11 (1996) 4173.
14. R. Forty, talk given at the Second International Conference on *B* Physics and *CP* Violation, Honolulu, Hawaii, March 24 - 27, 1997.
15. G.C. Branco, L. Lavoura, and F. Mota, *Phys. Rev.* D39 (1989) 3443.
16. S. Dimopoulos, L.J. Hall, and S. Raby, *Phys. Rev. Lett.* 68 (1992) 1984; Phys. R. Barbieri, L.J. Hall, and A. Romanino, *Phys. Lett.* B401 (1997) 47.
17. H. Fritzsch and X. Xing, *Phys. Rev.* D57 (1998) 594–597
18. A. Rasin, Report No. hep-ph/9708216.
19. Particle Data Group, R.M. Barnett *et al.*, *Phys. Rev.* D54 (1996) 1.
20. H. Fritzsch, *Phys. Lett.* B70 (1977) 436; B73 (1978) 317.
21. A.B. Carter and A.I. Sanda, *Phys. Rev. Lett.* 45 (1980) 952; I.I. Bigi and A.I. Sanda, *Nucl. Phys.* B193 (1981) 85.
22. A.J. Buras, *Report No. MPI-PhT/96-111*; and references therein.
23. L. Wolfenstein, *Phys. Rev. Lett.* 51 (1983) 1945.
24. See, e.g., K.S. Babu and Q. Shafi, *Phys. Rev.* D47 (1993) 5004; and references therein.
25. H. Fritzsch and Z. Xing, in preparation.